

The littlest Higgs model with T-parity and single top production at ep collision

Chong-Xing Yue, Jia Wen, Jin-Yan Liu, Wei Liu

Department of Physics, Liaoning Normal University, Dalian 116029, P.R.China *

March 11, 2008

Abstract

Based on calculating the contributions of the littlest Higgs model with T-parity (called LHT model) to the anomalous top coupling $tq\gamma$ ($q = u$ or c), we consider single top production via the t-channel partonic process $eq \rightarrow et$ at ep collision. Our numerical results show that the production cross section in the LHT model can be significantly enhanced relative to that in the standard model.

*E-mail:cxyue@lnnu.edu.cn

I. Introduction

The top quark with a mass of the order of the electroweak scale $m_t \sim 172\text{GeV}$ [1] is the heaviest particle yet discovered and might be the first place in which the new physics effects could be appeared. The correction effects of new physics on observables for top quark are often important than for other fermions. In particular, the anomalous top couplings, which affect top production and decay at high energy colliders, offer a unique place for testing the standard model (SM) flavor structure[2].

In the SM , the anomalous top quark couplings tqV ($q = c$ or u quark and $V = \gamma, Z$ or g gauge bosons), which are arised from the flavor changing (FC) interactions, vanish at tree level but can be generated at the one-loop level. However, they are very suppressed by the GIM mechanism, which can not be detected in the present and near future high energy experiments. It is well known that the anomalous top quark couplings tqV may be large in some new physics models beyond the SM and single top production is sensitive to these types of couplings. Thus, studying the contributions of the couplings tqV to single production of the top quark is of special interest. It will be helpful to test the SM flavor structure and new physics beyond the SM .

The ep collider, called $HERA$ collider with the center-of-mass ($c.m.$) energy $\sqrt{s} = 320\text{GeV}$ or $THERA$ collider with the $c.m.$ energy $\sqrt{s} = 1\text{TeV}$ [3], is the experimental facility where high energy electron-proton and positron-proton interactions can be studied. Within the SM , single top quark can not be produced at an observable level in the $HERA$ and $THERA$ collider experiments[4]. However, the $HERA$ and $THERA$ colliders can provide a good sensitivity on the couplings tqV via single top production[5]. Some studies about this type of single top quark production have appeared in the literature[6,7], which have shown that the $HERA$ and $THERA$ colliders are powerful tools for searching for the anomalous top quark couplings tqV .

The littlest Higgs model with T-parity (called LHT model)[8] is one of the attractive little Higgs models. To simultaneously implement T-parity, the LHT model introduces new mirror fermions. Under T-parity, particle fields are divided into T-even and T-odd sectors. The T-even sector consists of the SM particles and a heavy top quark T_+ , while

the T-odd sector contains heavy gauge bosons ($B_H, Z_H, W_H^\pm$), a scalar triplet (Φ), and so-called mirror fermions. The flavor mixing in the mirror fermion sector gives rise to a new source of flavor violation, which might generate significantly contributions to some flavor violation processes[9,10,11,12,13,14]. In this paper, we will concentrate our attention on single top production via the t-channel partonic process $eq \rightarrow et$ at the *HERA* and *THERA* collider experiments.

In the following section, we will give our numerical results in detail. Our conclusion and a simple discussion are given in section III.

II. Numerical results

At the *HERA* and *THERA* colliders, single top quark can be produced via the charged current (*CC*) process and the neutral current (*NC*) process. In the *SM*, the former process can proceed at the tree level, and its cross section is less than $1fb$ [15], while the latter process can only proceed at the one-loop level, which is *GIM* suppressed. However, the *NC* process $ep \rightarrow et + X$ is sensitive to the anomalous top quark couplings tqV .

The t-channel partonic process $eq \rightarrow et$ for the *NC* process $ep \rightarrow et + X$ can obtain contributions from the anomalous top quark couplings $tq\gamma$ and tqZ via γ exchange and Z exchange, respectively. However, the contribution from Z exchange is several orders of magnitude smaller than that from γ exchange. Thus, we will neglect the contributions of the coupling tqZ in our numerical estimation. Then, in the context of the *LHT* model, the differential cross section of the partonic process $e(P_e) + q(P_q) \rightarrow e(P_{e'}) + t(P_t)$ can be written as[5,6]:

$$\frac{d\hat{\sigma}(\hat{s})}{dt} = \frac{(K_{tq\gamma})^2 e^4}{32\pi \hat{s}^2} \left[\frac{-2\hat{s}^2 + 2\hat{s}m_t^2 - m_t^4}{t} + 2m_t^2 - 2\hat{s} \right], \quad (1)$$

where $t = q^2 = (P_{e'} - P_e)^2$ and $\sqrt{\hat{s}}$ is the *c.m.* energy of the t-channel partonic process $eq \rightarrow et$. $K_{tq\gamma}$ is the effective coupling constant of the $tq\gamma$ vertex contributed by the *LHT* model, in which we have drawn out one factor e . In above equation, we have neglected the masses of the incoming quarks and the electrons. However, for the Mandelstam variable t , to avoid divergence we will use $m_e = 0.511MeV$ for its upper and lower limits in the

phase space integral.




Figure 1: Feynman diagrams for the $tq\gamma$ ($q = u$, or c) vertex in the LHT model.

The effective cross section $\sigma(s)$ of single top production via the partonic process $eq \rightarrow et$ at ep collision can be obtained by folding the cross section $\hat{\sigma}(\hat{s})$ with the parton distribution function (PDF):

$$\sigma(s) = \sum_{q=u,c} \int_{x_{min}}^1 f_q(x, \mu) dx \int_{t_{min}}^{t_{max}} \frac{d\hat{\sigma}(\hat{s})}{dt} dt \quad (2)$$

with $x_{min} = \frac{m_t^2 + m_e^2}{s}$ and $\hat{s} = xs$, in which the $c.m.$ energy $\sqrt{s}$ is taken as $320 GeV$ for the $HERA$ collider and as $1 TeV$ for the $THERA$ collider. In our numerical calculation,

we will use *CTEQ6L PDFs*[16] for the quark distribution functions and assume that the factorization scale μ is of order m_t . The upper and lower limits of the Mandelstam variable t are taken as:

$$t_{max} \simeq -\frac{m_e m_t^2}{\hat{s}}, \quad t_{min} \simeq m_t^2 - \hat{s} - \frac{m_e^2(m_e^2 - m_t^2)}{\hat{s}}. \quad (3)$$

In the *LHT* model, the anomalous top quark coupling $tq\gamma$ can be induced by the interactions between the *SM* quarks and the mirror quarks mediated by T-odd gauge bosons ($B_H, Z_H, W_H^\pm$), as shown in *Fig.1*. The heavy scalar triplet Φ has not contributions to the coupling $tq\gamma$ at the order of ν^2/f^2 [10,11]. So, in our numerical estimation, we will neglect its contributions.

Using the relevant Feynman rules given in Refs.[10,17], we can calculate the contributions of the *LHT* model to the coupling constant $K_{tq\gamma}$ of the anomalous top quark coupling vertex $tq\gamma$. [Using the value of $K_{tq\gamma}$ for $q = c$, we have calculated the branching ratio $Br(t \rightarrow c\gamma)$ and find that our numerical result is approximately consistent with that given by Ref.[13]]. Observably, the cross section $\sigma(s)$ of single top production at *ep* collision is dependant on the model dependant parameters $f, M_{u_H^i}, M_{d_H^i}$, and $(V_{Hu})_{ij}$. The matrix elements $(V_{Hu})_{ij}$ can be determined via $V_{Hu} = V_{Hd}V_{CKM}^+$. The matrix V_{Hd} can be parameterized in terms of three mixing angles and three phases, which can be probed by *FCNC* processes in *K* and *B* meson systems, as discussed in detail in Refs.[10,12]. It is convenient to consider several representative scenarios for the structure of the matrix V_{Hd} . To simplify our calculation, we concentrate our study on the following two scenarios for the structure of the mixing matrix V_{Hd} :

Case I: $V_{Hd} = I, V_{Hu} = V_{CKM}^+$;

Case II: $s_{23}^d = \frac{1}{\sqrt{2}}, s_{12}^d = 0, s_{13}^d = 0, \delta_{12}^d = 0, \delta_{23}^d = 0, \delta_{13}^d = 0$.

It has been shown that, in both above cases, the constraints on the mass spectrum of the mirror fermions are very relaxed[10,12]. Furthermore, the masses of up and down type mirror quarks are equal to each other at the leading order of ν/f . Thus, we assume that the masses of mirror quarks have the relations $M_{u_H^i} = M_{d_H^i} = M_1 (i = 1, 2)$ and $M_{u_H^3} = M_{d_H^3} = M_2$. In our numerical estimation, we will take the scale parameter f and

the mass parameters M_1 , M_2 as free parameters.




Figure 2: In Case I, the cross section $\sigma(s)$ of single top production at the *THERA* collider with $\sqrt{s} = 1 \text{ TeV}$ as a function of f for different values of M_1 and M_2 .

Our numerical results for Case I are summarized in *Fig.2* and *Fig.3*. In these two figures, we have taken the values of the *CKM* matrix elements $(V_{CKM})_{ij}$ given by Ref.[18], in which V_{CKM} is constructed based on *PDG* parameterization[19]. For the *HERA* collider with $\sqrt{s} = 320 \text{ GeV}$, the value of $\sigma(s)$ is smaller than $1 \times 10^{-4} \text{ fb}$ in most of the parameter space of the *LHT* model. So, in *Fig.2*, we only plot the production cross section $\sigma(s)$ at the *THERA* collider with $\sqrt{s} = 1 \text{ TeV}$ as a function of the scale parameter f for different values of the mass parameters M_1 and M_2 . One can see from *Fig.2* that the value of the production cross section $\sigma(s)$ increases as the scale parameter

f decreases. The value of $\sigma(s)$ is insensitive to the mirror quark masses. The value of $\sigma(s)$ can reach $2.2 \times 10^{-3} fb$ for $f = 500 GeV$, $M_1 = 1500 GeV$ and $M_2 = 4000 GeV$, which is too small to be detected at the *THEIA* collider with $\sqrt{s} = 1 TeV$. In *Fig.3*, we plot the production cross section $\sigma(s)$ as function of the *c.m.* energy $\sqrt{s}$ for $M_1 = 1000 GeV$, $M_2 = 3000 GeV$, and three values of the scale parameter f . From *Fig.3*, we can see that, even for $f = 500 GeV$ and $\sqrt{s} = 1.5 TeV$, the value of $\sigma(s)$ is only $3.7 \times 10^{-3} fb$.




Figure 3: In Case I, the cross section $\sigma(s)$ a function of the *c.m.* energy $\sqrt{s}$ for $M_1 = 1000 GeV$, $M_2 = 3000 GeV$ and three values of f .

For Case I we have assumed $V_{Hd} = I$ and $V_{Hu} = V_{CKM}^+$, for Case II we will assume $V_{Hu} = V_{Hd} V_{CKM}^+$ and

$$V_{Hd} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ 0 & -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix}. \quad (4)$$

Observably, the value of the factor $\lambda_i = (V_{Hu})_{ij}^* (V_{Hu})_{i3}$ for Case II is different from that for Case I, which makes the effective production cross section $\sigma(s)$ of the t-channel partonic process $eq \rightarrow et$ for Case II differ from that for Case I. In *Fig.4*, the dependance of the cross section $\sigma(s)$ on the scale parameter f is presented for Case II. One can see




Figure 4: In Case II, the cross section $\sigma(s)$ at the *HERA* collider with $\sqrt{s} = 1TeV$ as a function of f for $M_2 = 3000GeV$ and three values of M_1 .

from *Fig.4* that the cross section $\sigma(s)$ of single top production for Case II is larger than that for Case I. For $f = 500GeV$, $M_2 = 3000GeV$, and $500GeV \leq M_1 \leq 1500GeV$, the value of $\sigma(s)$ is in the range of $5.9 \times 10^{-2}fb \sim 0.14fb$. In *Fig.5*, we plot $\sigma(s)$ as a function of the *c.m.* energy $\sqrt{s}$ for $M_1 = 1000GeV$, $M_2 = 3000GeV$ and three values of the scale parameter f . The value of $\sigma(s)$ can reach $0.18fb$ for $f = 500GeV$ and $\sqrt{s} = 1.5TeV$. If we assume that the *HERA* collider with $\sqrt{s} = 1TeV$ has a yearly integrated luminosity of $\mathcal{L} = 100fb^{-1}$, then there will be several tens of single top events to be generated.




Figure 5: Same as *Fig.3* but for Case II.

III. Conclusion and discussion

The *LHT* model is one of the attractive little Higgs models that is consistent with electroweak precision tests but also has a much richer flavor structure described by new flavor mixing matrices. This feature makes that the *LHT* model might generate significant contributions to some flavor violation processes. Based on calculating the contributions of the *LHT* model to the anomalous top quark coupling $tq\gamma$ ($q = u$ or c), single top quark production via the t-channel partonic process $eq \rightarrow et$ at ep collision is considered in this paper. Our numerical results show that the production cross section is too small to be measured at the *HERA* collider experiments. However, with reasonable values of the parameters in the *LHT* model, the production cross section can reach $0.14fb$ at the *THEIA* collider with $\sqrt{s} = 1TeV$. Certainly, the effects of the *LHT* model on single top quark production whether can be detected in future *THEIA* collider, depending on its luminosity.

Single top quark production at ep collision can also be induced by the anomalous top quark coupling tqg . Using the constraint on the single top production cross section obtained at the *HERA* collider, Ref.[20] has given upper limits on the coupling constant of the anomalous coupling tqg . The *LHT* model can generate a large anomalous top quark coupling tqg [13], which can also produce significant contributions to single top production at ep collisions. However, compared single top production based on the t-channel partonic process $eq \rightarrow et$, it has an additional light jet from a gluon or up type quarks. Thus, single top production at ep collision induced by the anomalous top quark coupling tqg should be further studied.

Acknowledgments

This work was supported in part by the National Natural Science Foundation of China under Grants No.10675057 and Foundation of Liaoning Educational Committee(2007T086).

References

- [1] CDF/DO Collab., <http://www-cdf.fnal.gov/physics/new/top/top.html>.
- [2] M. Benekce et al., *hep-ph/0003033*; D. Chakraborty, J. Konigsherg and D. Rainwater, *Ann. Rev. Nucl. Part. Sci.* **53**(2003)301; W. Wagner, *Rept. Prog. Phys.* **68**(2005)2409.
- [3] H. Abramowicz et al.[*TESLA-N Study Group Collaboration*], *Technical Design Report*, DESY 01-011 preprint; A. K. Cifci, S. Su Handoy, and Ö Yavas, *in Proc. of EPAC2000* P. 388; M. Klein, *THERA-electron-proton scattering at $\sqrt{s} = 1\text{TeV}$* , talk given at DIS'2000, Liverpool, April 25-30, 2000; P. J. Bussey, *Int. J. Mod. Phys. A* **17** (2002)1065.
- [4] U. Baur, J.J. Van der Bij, *Nucl. Phys. B* **304**(1988)451.
- [5] A. Belyaev, N. Kidonakis, *Phys. Rev. D* **65**(2002)037501.
- [6] H. Frizsch, D. Holtmannspötter, *Phys. Lett. B* **457**(1999)186; O. Cakir, S. Sultansoy, M. Yilmaz, *hep-ph/0105130*; A.T. Alan, A. Senol, *Europhys. Lett.* **57**(2002)669; N. Kidonakis and A. Belyaov, *JHEP* **0312**(2003)004.
- [7] W. Hollik, A. Hüttmann, B.A. Kniel, *Nucl. Phys. B* **790**(2008)200.
- [8] H. C. Cheng, I. Low, *JHEP* **0309**(2003)051; *JHEP* **0408**(2004)061; I. Low, *JHEP* **0410**(2004)067.
- [9] J. Hubisz, P. Meade, A. Noble, M. Perelstein, *JHEP* **0601**(2006)135.
- [10] M. Blanke et al., *JHEP* **0701**(2007)066.
- [11] A. Freitas, D. Wyler, *JHEP* **0611**(2006)061; M. Blanke et al., *JHEP* **0612**(2006)003.
- [12] J. Hubisz, S. J. Lee, G. Paz, *JHEP* **0606**(2006)041; M. Blanke et al., *Phys. Lett. B* **646**(2007)253.

- [13] Hou Hong-Sheng, *Phys. Rev. D* **75**(2007)094010.
- [14] M. Blanke et al., *Phys. Lett. B* **657**(2007)81; M. Blanke et al., *JHEP* **0706**(2007)082.
- [15] T. Stelzer, Z. Sullivan, S. Willenbrock, *Phys. Rev. D* **56**(1997)5919; S. Moretti, K. Odagivi, *Phys. Rev. D* **57**(1998)3040.
- [16] J. Pumplin et al., *JHEP* **0207**(2002)012; D. Stump et al., *JHEP* **0310**(2003)046.
- [17] J. Hubisz, P. Meade, *Phys. Rev. D* **71**(2005)035016.
- [18] G. Ahuja, M. Gupta, M. Randhawa, *hep-ph/0611324*.
- [19] W. -M. Yao et al., *J. Phys. G* **33**(2006)1.
- [20] A.A. Ashimva, S.R. Slabospitsky, *hep-ph/0604119*.